

Rochester Institute of Technology

RIT Digital Institutional Repository

Articles

Faculty & Staff Scholarship

2001

The Ramsey multiplicity of K_4

Konrad Piwakowski

Stanislaw Radziszowski

Follow this and additional works at: <https://repository.rit.edu/article>

Recommended Citation

Ars Combinatoria LX (2001) 131-136

This Article is brought to you for free and open access by the RIT Libraries. For more information, please contact repository@rit.edu.

The Ramsey Multiplicity of K_4

Konrad Piwakowski*

Department of Foundations of Informatics
Technical University of Gdańsk
80-952 Gdańsk, Poland
coni@pg.gda.pl

Stanisław P. Radziszowski

Department of Computer Science
Rochester Institute of Technology
Rochester, NY 14623, USA
spr@cs.rit.edu

Abstract

With the help of computer algorithms, we improve the lower bound on the Ramsey multiplicity of K_4 , and thus show that the exact value of it is equal to 9.

The Ramsey multiplicity $M(G)$ of a graph G is defined as the smallest number of monochromatic copies of G in any two-coloring of edges of $K_{R(G)}$, where $R(G)$ is the Ramsey number of G , i.e. the smallest integer n such that any two-coloring of edges of K_n contains monochromatic copy of G .

The study of Ramsey multiplicity was initiated in 1974 by Harary and Prins [3] who determined $M(G)$ for all graphs G of order four or less, except for K_4 and $K_4 - e$. The value of $M(K_4 - e)$ was later determined by Schwenk (cited in [2]). The upper bound $M(K_4) \leq 12$ was given in 1980 by Jacobson [4], and in 1988 Exoo [1] improved it by 3. The only nontrivial lower bound $M(K_4) \geq 4$ was recently presented by Olpp [7]. In this paper we improve this lower bound and thus show that $M(K_4) = 9$.

In the sequel, any two-coloring of the edges of K_n containing k monochromatic copies of K_4 is called an (n, k) -coloring. We say that two colorings are isomorphic if the graphs induced by the edges in the first color are isomorphic. Define $\mathcal{M}(n, k)$ to be set of all (n, k) -colorings. For a given (n, k) -coloring C let $H(C)$ denote the hypergraph formed by monochromatic copies of K_4 in C . Let us define $\mathcal{M}_d(n, k)$ to be the subset of all colorings $C \in \mathcal{M}(n, k)$ such that the maximal vertex degree in $H(C)$ is equal to d .

Our computational approach was to generate all nonisomorphic $(18, k)$ -colorings for $4 \leq k \leq 8$, by iterating an exhaustive enumeration of all possible one vertex extensions of $(n - 1, k - m)$ -colorings to (n, k) -colorings, for $m \geq 0$. Let us define $\mathcal{E}(n - 1, k - m, m)$ to be the subset of all colorings from $\mathcal{M}(n, k)$ which are one vertex extensions of some coloring from $\mathcal{M}(n - 1, k - m)$.

* Supported by the State Committee for Scientific Research KBN and academic computer center TASK.

Let $V(C)$ denote the set of vertices of coloring C . For each subset $W \subseteq V(C)$ let $N_3(W)$ denote the sum of the number of triangles in the first color induced by W in C and the number of triangles in the second color induced by $V(C) \setminus W$ in C . The following algorithm was used to perform the exhaustive search for all one vertex extensions $\mathcal{E}(n-1, k-m, m)$:

Algorithm 1

- Step 1: Initialize output set $Out = \emptyset$.
- Step 2: For each coloring C from $\mathcal{M}(n-1, k-m)$ execute steps 3, 4, 5.
- Step 3: For each subset $W \subseteq V(C)$ such that $N_3(W) = m$ execute steps 4, 5.
- Step 4: Create copy D of coloring C .
- Step 5: Add a new vertex v to coloring D . For each w in $V(C)$, assign color 1 to edge $\{v, w\}$, if $w \in W$, and assign color 2 to edge $\{v, w\}$, if $w \in V(C) \setminus W$. Add this coloring to Out .
- Step 6: Remove isomorphic copies from Out .

The following lemmas describe computational steps we followed in order to generate colorings of higher orders. As the initial step, we generated the set $\mathcal{M}(11, 0)$ by filtering out $(11, 0)$ colorings from all nonisomorphic graphs of order 11 (which were treated as two-colorings of K_{11}). The proofs of the lemmas are straightforward by considering degree sequences of all possible hypergraphs $H(C)$ in each case.

Lemma 1

$$\begin{aligned} \mathcal{M}(n, 0) &= \mathcal{E}(n-1, 0, 0), \quad \text{for } n \geq 2, \\ \mathcal{M}(n, k) &= \bigcup_{j=0}^{k-1} \mathcal{E}(n-1, j, k-j), \quad \text{for } k \geq 1, \text{ and } n \geq 2, \\ \mathcal{M}(16, 4) \setminus \mathcal{M}_1(16, 4) &= \bigcup_{j=0}^2 \mathcal{E}(15, j, 4-j). \end{aligned}$$

All the sets $\mathcal{M}(n, k)$, for $12 \leq n \leq 16$, and $0 \leq k \leq 3$ such that there is a nonempty entry for n, k in Table 1, were obtained by running Algorithm 1 for the terms on the right hand side of the first two rules in Lemma 1. For example, $\mathcal{M}(16, 3)$ was obtained by extending colorings from $\mathcal{M}(15, 0)$, $\mathcal{M}(15, 1)$ and $\mathcal{M}(15, 2)$.

The last identity in Lemma 1 describes the way of enumerating all $(16, 4)$ -colorings except those whose monochromatic copies of K_4 are vertex disjoint (denoted by $\mathcal{M}_1(16, 4)$). Unfortunately, there is a frightfully large number of $(13, 1)$ and $(14, 2)$ colorings, and we were not able to complete the sequence of extensions $\mathcal{M}(12, 0) \rightarrow \mathcal{M}(13, 1) \rightarrow \mathcal{M}(14, 2) \rightarrow \mathcal{M}(15, 3) \rightarrow \mathcal{M}_1(16, 4)$. Instead, in order to generate $\mathcal{M}_1(16, 4)$, we used the following approach:

Algorithm 2

Step 1: Generate the set of all 2-colorings of order 8 and extract from it $\mathcal{M}_1(8, 2)$.

Step 2: Generate $\mathcal{M}_1(12, 3)$ by exhaustively extending by 4 vertices all colorings in $\mathcal{M}_1(8, 2)$.

Step 3: Generate $\mathcal{M}_1(16, 4)$ by exhaustively extending by 4 vertices all colorings in $\mathcal{M}_1(12, 3)$.

In steps 2 and 3 exactly one new monochromatic K_4 is induced by 4 new vertices. As a result of the above algorithm we obtained 468 nonisomorphic $(16, 4)$ colorings.

The following lemma, together with Lemma 1, describes the remaining computational steps.

Lemma 2

$$\mathcal{M}(n, k) = \bigcup_{j=0}^{k-2} \mathcal{E}(n-1, j, k-j), \text{ for } k \geq 5, \text{ and } n \leq 19.$$

Using Algorithm 1 and Lemma 1 for $k \leq 4$, and Lemma 2 for $k \geq 5$, we were able to generate $\mathcal{M}(17, 0), \dots, \mathcal{M}(17, 6)$ and $\mathcal{M}(18, 0), \dots, \mathcal{M}(18, 8)$.

Table 1. The number of nonisomorphic (n, k) -colorings.

$n \setminus k$	0	1	2	3	4	5	6	7	8
11	546356								
12	1449166								
13	1184231								
14	130816	6144820							
15	640	50726	2491136						
16	2	28	382	19806	888440				
17	1	0	0	2	18	202	5757		
18	0	0	0	0	0	0	0	0	0

Table 1 presents the number of nonisomorphic (n, k) -colorings for all n and k , which were enumerated during our computations. The emptiness of the sets $\mathcal{M}(18, 0), \dots, \mathcal{M}(18, 8)$ implies the main theorem:

Theorem 1 $M(K_4) = 9$.

It is a natural goal to enumerate the set $\mathcal{M}(18, 9)$. Continuing our approach would require obtaining the whole set of colorings $\mathcal{M}(17, 7)$. The latter was unfeasible, and we were able to enumerate only the set $\mathcal{M}(17, 6)$.




Figure 1. The new $(18, 9)$ -coloring

Since, similar to the previous lemmas, we easily have

$$\mathcal{M}(18, 9) \setminus \mathcal{M}_2(18, 9) = \bigcup_{j=0}^6 \mathcal{E}(17, j, 9 - j),$$

we enumerated all $(18, 9)$ -colorings such that not every vertex belongs to exactly two monochromatic copies of K_4 . There are 4 such colorings, where two of them come from the other two by exchanging the colors. Of the two essentially different colorings, one was presented in [1] and the other is presented in Figure 1, where only the edges in one color are shown. There are seven K_4 in the first color induced by vertex sets: $\{A, B, D, E\}$, $\{B, C, E, F\}$, $\{D, E, G, H\}$, $\{E, F, H, I\}$, $\{J, C, G, M\}$, $\{K, A, I, L\}$, $\{J, K, L, M\}$ and two K_4 in the second color induced by $\{B, O, P, H\}$

and $\{N, D, F, Q\}$. Notice that the labels Q and R in the Figure 2 in [1] are mistakenly switched. It results in serious complications with decoding the $(18, 9)$ -coloring by the reader.

The question about contents of the set $\mathcal{M}_2(18, 9)$ remains open; however we conjecture that it is empty.

Three powerful programs, *nauty*, *makeg*, and *autoson*, implemented by Brendan McKay [5] were used in our work. All the algorithms specific for this project were written independently by both authors, and then a very large number of intermediate and final graphs were tested for isomorphism between the two implementations. Moreover, the cardinalities of all sets $\mathcal{M}(n, 0)$, for $n = 11, \dots, 18$ agreed with the previous enumeration described in [6].

References

- [1] G. Exoo, Some Constructions Related to Ramsey Multiplicity, *Ars. Comb.* **26** (1988) 233-242.
- [2] F. Harary, A survey of Generalized Ramsey Theory, in: Graphs and Combinatorics (R. Bari and F. Harary, eds.), *Lecture Notes in Mathematics* **406**, Springer (1974) 10-17.
- [3] F. Harary, G. Prins, Generalized Ramsey Theory for Graphs IV, the Ramsey Multiplicity of a Graph, *Networks* **4** (1974) 163-173.
- [4] M. S. Jacobson, A note on Ramsey multiplicity, *Discrete Math.* **29** (1980) 201-203.
- [5] B. D. McKay, *Nauty User's Guide* (version 1.5), Technical report TR-CS-90-02, Computer Science Department, Australian National University, 1990. WWW reference: <http://cs.anu.edu.au/people/bdm/>.
- [6] B. D. McKay, S. P. Radziszowski, A New Upper Bound for the Ramsey Number $R(5, 5)$, *Australasian Journal of Combinatorics*, **5** (1992), 13-20.
- [7] D. Olpp, A Lower Bound for the Ramsey Multiplicity of K_4 , *Ars. Comb.* **49** (1998) 289-295.